

THE ISOPERIMETRIC INEQUALITY FOR $\mathrm{SL}(n, \mathbb{Z})$

organized by

Nathan Broaddus, Tim Riley, and Kevin Wortman

Workshop Summary

1. PROGRESS AND PROGRAMS FOR SOLUTIONS

During the week of September 8-12, 2008 we held a workshop at AIM to answer the following question.

Does $\mathrm{SL}(n, \mathbb{Z})$ admit a quadratic isoperimetric inequality when $n \geq 4$?

We made progress in addressing this problem from the three perspectives below.

1.1. Modifying horoballs. A standard approach to constructing a quasi-isometric model for $\mathrm{SL}(4, \mathbb{Z})$ is to start with the symmetric space $X = \mathrm{SL}(4, \mathbb{R})/\mathrm{SO}(4, \mathbb{R})$ and remove a union of open horoballs to get a space

$$Y = X - \bigcup_{r \in \mathcal{R}} H_r$$

where \mathcal{R} is an $\mathrm{SL}(4, \mathbb{Z})$ -equivariant collection of geodesic rays in X . Here each horoball H_r is constructed using a Buseman function corresponding to the ray $r \in \mathcal{R}$. Since X is nonpositively curved it satisfies a quadratic isoperimetric inequality. Therefore, given a loop in Y one may fill it with a disk $D \subset X$ whose area is quadratic in the length of the loop.

One would then like to push D out of each horoball H_r while fixing ∂D and increasing its area in a controlled (hopefully Lipschitz) manner. One difficulty with this approach is that one may find arbitrarily large sets of horoballs $\{H_1, \dots, H_k\}$ with nonempty intersection

$$\bigcap_{i=1}^k H_i \neq \emptyset.$$

We considered an alternative collection \mathcal{U} of subsets X with finite intersection properties, and such that

$$Y = X - \bigcup_{H \in \mathcal{U}} H$$

For the rings $A = \mathbb{Z}, \mathbb{R}$ Let N be the unipotent group

$$N(A) = \left\{ \begin{pmatrix} 1 & a_{11} & \cdots & a_{1n} \\ 0 & 1 & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix} \mid a_{ij} \in A \right\}$$

The quotient $\mathrm{SL}(4, \mathbb{Z}) \backslash X$ has a thick/thin decomposition where the noncompact thin part is isometric to a “semidirect product” $S \ltimes (N(\mathbb{Z}) \backslash N(\mathbb{R}))$ where S is the Euclidean sector (and commutative monoid)

$$S = \left\{ \begin{pmatrix} e^{t_1} & 0 & 0 & 0 \\ 0 & e^{t_2} & 0 & 0 \\ 0 & 0 & e^{t_3} & 0 \\ 0 & 0 & 0 & e^{t_4} \end{pmatrix} \mid t_i \in \mathbb{R}, \sum_{i=1}^n t_i = 0, t_1 \geq \dots \geq t_4 \right\}.$$

Given $i, j \in \{1, 2, 3, 4\}$, we let $S_{i=j}$ be all the points in S with $t_i = t_j$. For any subset $S' \subseteq S$, we let $F(S')$ be a fundamental domain in X for $S' \ltimes (N(\mathbb{Z}) \backslash N(\mathbb{R}))$, and then we let $P(S') \leq \mathrm{SL}(4, \mathbb{Z})$ be the group that stabilizes $F(S')(\infty)$ and take $H(S') = P(S')F(S')$. Our set \mathcal{U} of modified horoballs is the set of $\mathrm{SL}(4, \mathbb{Z})$ translates of $H(S)$, $H(S_{1=2})$, $H(S_{2=3})$, $H(S_{3=4})$, $H(S_{1=2=3})$, $H(S_{2=3=4})$, and $H(S_{1=2,3=4})$.

Now to prove that $\mathrm{SL}(4, \mathbb{Z})$ has a quadratic Dehn function, one has to show that the boundaries of each of the first four of modified horoballs listed above satisfies a quadratic Dehn function. With that knowledge, we can assume that our disc $D \subseteq X$ can be replaced by a disc with the same boundary, with roughly the same area, that lies either in the thick part of X or else in one of the last three types of modified horoballs listed above. Having a disc entirely in the thick part is our goal, so the next step is to analyze the geometry of the last three types of modified horoballs. These horoballs reflect the geometry of groups like $\mathrm{SL}(3, \mathbb{Z}) \ltimes \mathbb{Z}^3$.

1.2. Recursive combinatorial subdivision of spanning disks. A number of combinatorial ideas were explored. A suggestion of Kassabov is to consider a weighted infinite presentation of the Steinberg group $\mathrm{St}(4, \mathbb{Z})$ which is quasi-isometric to $\mathrm{SL}(4, \mathbb{Z})$. Then Steinberg group has an infinite presentation with generators $e_{ij}(\lambda)$ where $1 \leq i \neq j \leq 4$ and $\lambda \in \mathbb{Z}$ relations meant to model the relations behavior of elementary matrices:

$$e_{ij}(\lambda)e_{ij}(\nu) = e_{ij}(\lambda + \nu) \tag{1}$$

$$[e_{ij}(\lambda), e_{kl}(\nu)] = 1 \tag{2}$$

$$[e_{ij}(\lambda), e_{jk}(\nu)] = e_{kl}(\lambda\nu) \tag{3}$$

Instead of assigning length 1 and area 1 to each generator and relation one should consider the length of $e_{ij}(\lambda)$ to be $K_1 \log(\lambda)$, the triangular relations of type (1) should have area $K_2 \log(\lambda) \log(\nu)$, the quadrilateral relations of type (2) should have area $K_3 \log(\lambda) \log(\nu)$, and the pentagon relations of type (3) should have area $K_4 \log(\lambda) \log(\nu)$. One may then try to prove that with this weighted presentation $\mathrm{SL}(4, \mathbb{Z})$ has a quadratic isoperimetric inequality and that this weighted infinite presentation give the same isoperimetric inequality as a finite presentation up to linear constants.

An argument of this type shows great promise of answering the a related conjecture that $\mathrm{SL}(3, \mathbb{Z})$ has linear filling length.

1.3. Homotopy type of the asymptotic cone. A nonconstructive approach to the problem is to try to describe the homotopy type of the asymptotic cone of $\mathrm{SL}(4, \mathbb{Z})$. Simple connectivity of every asymptotic cone of $\mathrm{SL}(4, \mathbb{Z})$ would imply a quadratic Dehn function [Riley2003].

The asymptotic cone $X = \text{Cone}_\infty(\text{SL}(4, \mathbb{R}))$ for $\text{SL}(4, \mathbb{R})$ is a Euclidean \mathbb{R} -building that one gets by a construction of Bruhat and Tits from considering a (nondiscrete) valuation on the Robinson field which is a quotient of a subring of the hyperreal numbers [KSTT]. Since $\text{SL}(4, \mathbb{Z})$ is quasi-isometrically embedded in $\text{SL}(4, \mathbb{R})$ one may view $\text{Cone}_\infty(\text{SL}(4, \mathbb{Z}))$ to be a subset of X . In fact, one may describe $\text{Cone}_\infty(\text{SL}(4, \mathbb{Z}))$ as a complement of a set of horoballs \mathcal{H} in X .

The Euclidean \mathbb{R} -building X us a union of subsets $A \in \mathcal{A}$ isometric to Euclidean space \mathbb{R}^{4-1} called apartments. If one can identify the homotopy type of

$$A - \bigcap_{H \in \mathcal{H}} H$$

for every apartment $A \in \mathcal{A}$ and further understand the homotopy types of intersections of such sets then one might be able to show simple connectivity for $\text{Cone}_\infty(\text{SL}(4, \mathbb{Z}))$ giving a polynomial Dehn function.

Bibliography

- [Drutu2004] C. Drutu. Filling in solvable groups and in lattices in semisimple groups. *Topology*, 43:983–1033, 2004.
- [Epstein1992] D. B. A. Epstein, J. W. Cannon, D. F. Holt, S. V. F. Levy, M. S. Paterson, and W. P. Thurston. *Word Processing in Groups*. Jones and Bartlett, 1992.
- [Gersten1993] S. M. Gersten. Isoperimetric and isodiametric functions. In G. Niblo and M. Roller, editors, *Geometric group theory I*, number 181 in LMS lecture notes. Camb. Univ. Press, 1993.
- [Gromov1993] M. Gromov. Asymptotic invariants of infinite groups. In G. Niblo and M. Roller, editors, *Geometric group theory II*, number 182 in LMS lecture notes. Camb. Univ. Press, 1993.
- [KSTT] L. Kramer et al., Asymptotic cones of finitely presented groups. *Adv. Math.*, 193(1):142–173, 2005.
- [LeuzingerPittet2004] E. Leuzinger and Ch. Pittet. On quadratic Dehn functions. *Math. Z.*, 248(4):725–755, 2004.
- [LMR1993] A. Lubotzky, S. Mozes, and M. S. Raghunathan. Cyclic subgroups of exponential growth and metrics on discrete groups. *C.R. Acad. Sci. Paris, Série 1*, 317:723–740, 1993.
- [LMR2000] A. Lubotzky, S. Mozes, and M. S. Raghunathan. The word and Riemannian metrics on lattices of semisimple groups. *Inst. Hautes Études Sci. Publ. Math.*, 91:5–53, 2000.
- [Milnor1971] J. Milnor. *Introduction to algebraic K-theory*, volume 72 of *Annals of Mathematical Studies*. Princeton University Press, 1971.
- [Riley2005] T. R. Riley. Navigating the Cayley graphs of $\text{SL}_N(\mathbb{Z})$ and $\text{SL}_N(\mathbb{F}_p)$. *Geometriae Dedicata*, 113(1):215–229, 2005.
- [Riley2003] T. R. Riley. Higher connectedness of asymptotic cones. *Topology*, 42(6):1289–1352, 2003.
- [Saper1997] L. Saper. Tilings and finite energy retractions of locally symmetric spaces, Comment. Math. Helv. **72** (1997), no. 2, 167–202.